

AD-A052 313

FLORIDA STATE UNIV TALLAHASSEE DEPT OF STATISTICS
ON THE STRONG LAW OF LARGE NUMBERS AND RELATED RESULTS FOR QUAS--ETC(U)
AUG 77 R J SERFLING

F/G 12/1

N00014-76-C-0608

UNCLASSIFIED

FSU-STATISTICS-M430

NL

[OF]

AD
A052313



END
DATE
FILMED

5-78

DDC

AD A 052313

AD No. _____
DDC FILE COPY

**The Florida State University
Department
of
Statistics
Tallahassee, Florida
32306**



(12)

ON THE STRONG LAW OF LARGE NUMBERS AND
RELATED RESULTS FOR QUASI-STATIONARY SEQUENCES

by R. J. Serfling

FSU Statistics Report M430
ONR Technical Report No. 123

August, 1977
Department of Statistics
The Florida State University
Tallahassee, Florida 32306

Research supported by the Office of Naval Research under Contract No.
N00014-76-C-0608. Reproduction in whole or in part is permitted for
any purpose of the United States Government.

DDC
RECEIVED
APR 7 1978
B

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

ABSTRACT

ON THE STRONG LAW OF LARGE NUMBERS AND RELATED RESULTS FOR QUASI-STATIONARY SEQUENCES

Under second moment assumptions and weak dependence conditions on a sequence of random variables $\{X_i\}$, Gaposkin (1975) has established almost sure convergence of the series $\sum_1^\infty \lambda_k X_k$ under certain restrictions on the rate of convergence to 0 of the constants $\{c_k\}$. Similarly, Móricz (1977) has established conditions for the almost sure convergence to 0 of the sequence $\lambda_n \sum_1^n X_k$. In the present paper, some extensions of these results are obtained.

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DDC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION _____		
BY _____		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL. and/or	SPECIAL
A		

1. Main results and discussion. Consider a sequence of random variables $\{X_i\}$ satisfying

$$EX_i \equiv 0, \quad EX_i^2 \equiv 1$$

and, for a sequence of constants $\{\phi_i\}$,

$$|EX_j X_k| \leq \phi_{k-j}, \text{ all } j \leq k.$$

Such a sequence $\{X_i\}$ is called *quasi-stationary* with respect to the sequence $\{\phi_i\}$. The almost sure asymptotic behavior of the sum $\sum_{i=1}^n X_i$ may be characterized by an assertion of the form

$$(A) \quad \lambda_n \sum_{i=1}^n X_i \rightarrow 0, \quad n \rightarrow \infty, \quad \text{w.p.1,}$$

where $\{\lambda_n\}$ is a sequence of positive constants tending to 0. It is of interest to establish (A) under mild restrictions on the constants $\{\phi_i\}$ and $\{\lambda_i\}$. A related problem concerns the almost sure behavior of the sum $\sum_{i=1}^n \lambda_i X_i$ for such a sequence of constants $\{\lambda_n\}$. In this case the desired assertion is

$$(\alpha) \quad \sum_{i=1}^{\infty} \lambda_i X_i \text{ converges w.p.1.}$$

By the well-known Kronecker lemma, (α) implies (A) in the case of λ_n non-increasing.

AMS 1970 *subject classifications*. Primary 60F99, Secondary 60G99.

Key words and phrases. Quasi-stationary random variables; strong law of large numbers; almost sure convergence of infinite series.

Rademacher (1922) and Mensov (1923) independently established that

(α) holds if

$$(1.1a) \quad \phi_1 = 0, \quad i > 0,$$

and

$$(1.1b) \quad \sum_1^\infty \lambda_n^2 \log^2 n < \infty.$$

Kac, Salem and Zygmund (1948) relaxed (1.1a) to $\phi_n = O(n^{-1-\epsilon})$ for an $\epsilon > 0$.

Gaposkin (1975) proved the following much broader result. Put

$$w(n) = \sum_{i=1}^n \phi_i.$$

THEOREM α (Gaposkin). *If*

$$(1.2) \quad \sum_1^\infty w(n) \lambda_n^2 \log^2 n < \infty,$$

then (α) holds.

This theorem allows the possibility of $w(n) \rightarrow \infty$, whereas the earlier results are confined to the case $w(\infty) < \infty$.

Returning to (A), we have

COROLLARY α . *If (1.2) is satisfied and λ_n is nonincreasing, then (A) holds.*

On the other hand, a direct approach — bypassing (α) — offers the possibility of obtaining (A) under weaker restrictions than (1.2). In this direction, Móricz (1977) has obtained the following result.

THEOREM A (Móricz). If

$$(1.3a) \quad \sum_1^{\infty} w(n) \lambda_n^2 < \infty,$$

$$(1.3b) \quad w(n) \lambda_n^2 \text{ is nonincreasing,}$$

and

$$(1.3c) \quad w(2n)/w(n) \geq q > 1, \text{ all } n,$$

then (A) holds.

Note that (1.3a) relaxes (1.2). However, (1.3c) requires $w(n)$ to grow at a fast rate. For example, (1.3c) is satisfied by $w(n)$ of the form $w(n) = cn^a$, but not by $w(n)$ of the form $w(n) = \exp(2\sqrt{\log n})$. For the latter, Theorem A is inapplicable, whereas Theorem α does yield a conclusion.

The present note provides an alternate to Theorem A which essentially removes condition (1.3c). As in [5], put $W(1) = w(1)$ and, for $n \geq 2$, define $W(n)$ by

$$W^h(n) = W^h([1/2]n) - 1 + w^h([1/2]n).$$

THEOREM B. If

$$(1.4a) \quad \sum_1^{\infty} W(n) \lambda_n^2 < \infty$$

and

(1.46) $W(n) \lambda_n^2$ is nonincreasing,

then (A) holds.

Conditions (1.3a) and (1.3c) together imply (1.4a), as evident from the Lemma below. Also, the mild constraints (1.3b) and (1.4b) are mere variants of each other. Thus Theorem B has somewhat broader application than Theorem A. In particular, it yields

EXAMPLE. Consider $w(n) = \exp(2\sqrt{\log n})$. In this case (by the Lemma below)

$$W(n) = O(w(n) \log n),$$

so that (A) holds if λ_n satisfies (1.4b) for this $w(n)$ and if

$$(1.5) \quad \sum_1^\infty w(n) \lambda_n^2 \log n < \infty. \quad \square$$

In the preceding example, the use of Corollary α would be less effective than Theorem B, since (1.5) is weaker than (1.2). The gain in effectiveness of Theorem B over Corollary α occurs when $w(n)$ grows sufficiently fast.

LEMMA. (i) In general, $W(n) = O(w(n) \log^2 n)$.

(ii) If $w(n) = \exp(2\sqrt{\log n})$, then $W(n) = O(w(n) \log n)$.

(iii) If $w(n)$ satisfies (1.5c), then $W(n) = O(w(n))$.

As a complement to Theorem B, the following generalization of Theorem α will be established.

THEOREM β . If (1.4a) and

$$(1.6a) \quad \sum_1^{\infty} w(n) \lambda_n^2 (\log n) (\log \log n)^{1+\epsilon} < \infty, \text{ for some } \epsilon > 0,$$

are satisfied, then (a) holds.

Since (1.2) implies each of (1.4a) and (1.6a), Theorem β generalizes Theorem α .

2. Proofs.

PROOF OF THE LEMMA. Note that, for $2^k \leq n < 2^{k+1}$,

$$(2.1) \quad W^{\frac{1}{2}}(n) < W^{\frac{1}{2}}(2^{k+1} - 1) = \sum_{j=0}^k w^{\frac{1}{2}}(2^j).$$

Thus $W^{\frac{1}{2}}(n) \leq k w^{\frac{1}{2}}(n) = O(w^{\frac{1}{2}}(n) \log n)$, which gives (i). Now, for $j \leq k$,

$$\begin{aligned} \exp \sqrt{j} &= \exp \sqrt{k} \exp \left(\frac{1-k}{\sqrt{j} + \sqrt{k}} \right) \\ &\leq \exp \sqrt{k} \exp \left(\frac{1-k}{2\sqrt{k}} \right) \\ &= \exp \frac{1}{2}\sqrt{k} \left(\exp \frac{1}{2\sqrt{k}} \right)^j. \end{aligned}$$

Thus by (2.1) we obtain, for the case $w(n) = \exp(2\sqrt{\log n})$, that

$$W^{\frac{1}{2}}(n) \leq (\exp \frac{1}{2}\sqrt{k}) \frac{\exp \sqrt{k}}{\exp \left(\frac{1}{2\sqrt{k}} \right) - 1} \leq 2\sqrt{k} \exp \sqrt{k},$$

i.e., $W^{\frac{1}{2}}(n) = O(\sqrt{\log n} w^{\frac{1}{2}}(n))$, so that (ii) is proved. Finally, for $w(n)$ satisfying (1.3c), the use of (2.1) yields

$$W^{\frac{1}{2}}(n) \leq w^{\frac{1}{2}}(2^k) \sum_{j=0}^k \left(\frac{1}{q}\right)^{\frac{1}{2}(k-j)},$$

from which (iii) follows. \square

In proving Theorems B and β , the following maximal inequality will be used.

LEMMA 2.1. For $m \geq 1$, $n \geq 1$,

$$E \left\{ \max_{1 \leq k \leq n} \left(\sum_{i=m+1}^{m+k} a_i X_i \right)^2 \right\} \leq 2W(n) \sum_{i=m+1}^{m+n} a_i^2.$$

This was proved by Móricz (1976), extending an earlier result of Serfling (1970a). For Theorem β , we will also need the following easily proved parallel result [5], [8].

LEMMA 2.2. For $n \geq 1$,

$$E \left\{ \left(\sum_{i=m+1}^{m+n} a_i X_i \right)^2 \right\} \leq 2w(n) \sum_{i=m+1}^{m+n} a_i^2.$$

PROOF OF THEOREM B. In order to show (A), it is equivalent to show that for every $\epsilon > 0$,

$$(2.2) \quad P\{|\lambda_n S_n| > \epsilon \text{ infinitely often}\} = 0.$$

Now observe that the nonincreasingness of $\lambda_n^2 W(n)$, combined with the non-decreasingness of $W(n)$, implies that λ_n is nondecreasing. Thus, by the Borel-Cantelli lemma, (2.2) holds if

$$(2.3) \quad \sum_{k=0}^{\infty} P\{\lambda_{2^k} \max_{2^k \leq n < 2^{k+1}} |S_n| > \epsilon\} < \infty.$$

Now

$$\begin{aligned} P\{\lambda_{2^k} \max_{2^k \leq n < 2^{k+1}} |S_n| > \epsilon\} \\ \leq \epsilon^{-2} \lambda_{2^k}^2 E\{\max_{2^k \leq n < 2^{k+1}} S_n^2\} \\ \leq 2\epsilon^{-2} \lambda_{2^k}^2 E\{S_{2^k}^2 + \max_{2^k \leq n < 2^{k+1}} (S_n - S_{2^k})^2\}. \end{aligned}$$

A two-fold application of Lemma 2.1 gives

$$P\{\lambda_{2^k} \max_{2^k \leq n < 2^{k+1}} |S_n| > \epsilon\} \leq 4\lambda_{2^k}^2 W(2^k)2^k.$$

Thus the sum in (2.3) is bounded by $4 \sum_{k=0}^{\infty} 2^k \lambda_{2^k}^2 W(2^k)$, which in turn

in view of (1.3b) is clearly bounded by $4 \sum_{n=1}^{\infty} \lambda_n^2 W(n)$. By (1.3a), the required (2.3) thus holds. \square

PROOF OF THEOREM β . Following the approach of [1], and using a standard elementary argument, we first establish that T_{2^k} converges w.p.1 to a limit T_{∞} , by showing that

$$(2.4) \quad \sum_{k=0}^{\infty} ||\Delta_k|| < \infty,$$

where $\Delta_k = S_{2^{k+1}} - S_{2^k}$ and $||\Delta_k||$ denotes $(E \Delta_k^2)^{\frac{1}{2}}$. By the Cauchy-

Schwarz inequality,

$$\sum ||\Delta_k|| \leq \left(\sum d_k^2 ||\Delta_k||^2 \right) \left(\sum d_k^{-2} \right)$$

for positive constants d_k . Choose $d_k = k^{\frac{1}{2}} (\log k)^{\frac{1}{2}(1+\epsilon)}$, $\epsilon > 0$. Then, applying Lemma 2.2 and (1.6a), we obtain for an appropriate constant C ,

$$\begin{aligned} \sum ||\Delta_k|| &\leq C \sum k (\log k)^{1+\epsilon} [2w(2^k) \sum_{j=2^{k+1}}^{2^{k+1}-1} \lambda_j^2] \\ &\leq 2C \sum_{n=1}^{\infty} (\log n) (\log \log n)^{1+\epsilon} w(n) \lambda_n^2 < \infty. \end{aligned}$$

Next we establish that T_n converges w.p.1 to T_{∞} , by showing that

$$\max_{2^k \leq n < 2^{k+1}} |T_n - T_{2^k}| \longrightarrow 0 \text{ w.p.1.}$$

This follows, by an argument similar to the proof of Theorem B, if

$$\sum_{k=0}^{\infty} E \left\{ \max_{2^k \leq n < 2^{k+1}} (T_n - T_{2^k})^2 \right\} < \infty,$$

which in turn is established by Lemma 2.2 and (1.4a), via

$$2 \sum_{k=0}^{\infty} w(2^k) \sum_{n=2^k}^{2^{k+1}-1} \lambda_n^2 \leq \sum_{n=1}^{\infty} w(n) \lambda_n^2 < \infty. \quad \square$$

REFERENCES

- [1] Gaposkin, V. F. (1975). Convergence of series which correspond to stationary sequences. (In Russian) *Izv. Akad. Nauk. SSSR, Ser. Mat.* 39 1366-1392.
- [2] Kac, M., Salem, R. and Zygmund, A. (1948). A gap theorem. *Trans. Amer. Math. Soc.* 63 235-248.
- [3] Mensov, D. (1923). Sur les séries de fonctions orthogonales, I. *Fund. Math.* 4 82-105.
- [4] Móricz, F. (1976). Moment inequalities and the strong laws of large numbers. *Zeitschrift für Wahrscheinlichkeitstheorie und verw. Gebiete* 35 299-314.
- [5] Móricz, F. (1977). The strong laws of large numbers for quasi-stationary sequences. *Ibid.* 38 223-236.
- [6] Rademacher, H. (1922). Einige Sätze über Reihen von allgemeinen Orthogonal-functionen. *Math. Ann.* 87 112-138.
- [7] Serfling, R. J. (1970a). Moment inequalities for the maximum cumulative sum. *Ann. Math. Statist.* 41 1227-1234.
- [8] Serfling, R. J. (1970b). Convergence properties of S_n under moment restrictions. *Ann. Math. Statist.* 41 1235-1248.

14 FSU-STATISTICS-M430
TR-123-ONR

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1. REPORT NUMBER FSU No. M430✓ ONR No. 123✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE 6 ON THE STRONG LAW OF LARGE NUMBERS AND RELATED RESULTS FOR QUASI-STATIONARY SEQUENCES.	5. TYPE OF REPORT & PERIOD COVERED 9 Technical Report.	
7. AUTHOR(s) 10 R. J. Serfling	6. PERFORMING ORGANIZATION REPORT NUMBER FSU Statistics Report M430	
9. PERFORMING ORGANIZATION NAME AND ADDRESS The Florida State University Department of Statistics Tallahassee, Florida 32306	8. CONTRACT OR GRANT NUMBER(s) 15 ONR No. N00014-76-C-0608✓	
11. CONTROLLING OFFICE NAME & ADDRESS Office of Naval Research Statistics & Probability Program Arlington, Virginia 22217	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 12/13p.	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE 11 Aug 1977	
	13. NUMBER OF PAGES 9	
	15. SECURITY CLASS (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS Quasi-stationary random variables; Strong law of large numbers; Almost sure convergence of infinite series.		

20. ABSTRACT

Under second moment assumptions and weak dependence conditions on a sequence of random variables $\{X_i\}$, Gaposkin (1975) has established almost sure convergence of the series $\sum_{k=1}^{\infty} \lambda_k X_k$ under certain restrictions on the rate of convergence to 0 of the constants $\{\lambda_k\}$. Similarly, Móricz (1977) has established conditions for the almost sure convergence to 0 of the sequence $\lambda_n \sum_{k=1}^n X_k$. In the present paper, some extensions of these results are obtained.

Handwritten notes:
 $\langle X_{sub\ 1} \rangle$
 $\langle c_{sub\ k} \rangle$
 $(\lambda_{sub\ n}) (\text{sum from 1 to } n \text{ of } X_{sub\ k})$
 $\sum_{k=1}^{\infty} \lambda_k X_k$
 $\sum_{k=1}^n \lambda_k X_k$
 400 277
 sum from 1 to infinity of $\lambda_{sub\ k} X_{sub\ k}$
 Ser